

# DETECTION OF HONEY BEE VIRUSES IN JORDAN

Nizar Haddad

Bee Research Unit & Biotechnology Unit. National Center for Agricultural Research and Extension.  
Baq'a 19381 Jordan, drnizarh@yahoo.com . www.jordanbru.info

---

## **ABSTRACT**

Mortality of honeybees is a serious problem that beekeepers have to face periodically in Jordan and worldwide. The presence of honeybee viruses, in addition to other pathogens and parasites may be one of its possible causes.

In this work, individual bee samples from the Ajlon area of Jordan were screened for six honey bee viruses using RT-PCR and specific primers. Acute bee paralysis virus (ABPV), Sacbrood virus (SBV) and Deformed wing virus (DWW) were detected in the samples, while Black queen cell virus (BQCV), Chronic bee paralysis virus (CBPV) and Kashmir bee virus (KBV) were not detected. This is the first report of the detection of bee viruses in Jordan utilizing molecular techniques.

Key words: Honeybee viruses, RT-PCR, molecular markers

## **INTRODUCTION**

The beekeeping industry plays a key role in agricultural production in Jordan. A recent study estimated that, in Jordan, the total production value of 12 bee-pollinated crops was \$117.4M in 2005, of which \$50.7M could be directly attributed to pollination by honey bees (Haddad et al., 2007), which is over 16 times the value of the locally produced honey (\$3.1M). Recently a large proportion of beekeepers complained of considerable bee mortality in their hives, especially in the Ajlon province. Mass mortality of honeybees (*Apis mellifera*) is one of the most serious problems that beekeepers and dependent industries face worldwide. A graphic example is the widely publicised economic, social and political upheaval caused by the extensive colony collapse among migratory beekeepers in the USA (Cox-Foster et al., 2007; Stokstad, 2007a; 2007b), and similar colony collapses and agricultural disruption during the previous century (Beuhne, 1910; Wilson and Menapace, 1979; Oertel, 1965; Foote, 1966; Rennie, 1921; Bailey, 1964; Goodacre, 1943).

There are many factors that may contribute to bee mortality, including bee parasites and diseases, (Ball and Bailey, 1997; Cox-Foster et al., 2007) the effects of etiological agents used to treat such diseases, climatic factors (Bailey et al., 2001; Wilson and Menapace, 1979) and exposure to insecticides or pesticides used in agriculture (Antunez et al., 2006; Suchail et al., 2004).

The association of viruses with honeybee mortality is a particular cause for concern world-wide (Bailey et al., 1981; 1983; Hornitzky, 1987; Allen and Ball, 1996; Nordstrom et al., 1999; Tentcheva et al., 2004a,b; Todd et al., 2007;). The honeybee is host to at least 18 viruses (Bailey and Ball, 1991). All are normally asymptomatic and persist in the colonies as covert infections (Bailey and Ball, 1991; Ball and Bailey, 1997). Some, such as Chronic bee paralysis virus (CBPV), Black queen cell virus (BQCV), Sacbrood virus (SBV), Deformed wing virus (DWW) and Cloudy wing virus (CWV), will also produce recognizable clinical symptoms at certain life stages (Bailey and Ball, 1991; Allen and Ball, 1996). Nucleotide sequences are now available for many bee viruses, enabling the development of nucleic acid based diagnostic assays (e.g. Tentcheva et al., 2004b; Chen et al., 2005; Topley et al., 2005; Chantawannakul et al., 2006; Berenyi et al., 2006; Grabesteiner et al., 2007). In this study we use several such assays to assess the presence of honeybee viruses in some Ajlon province apiaries.

## **MATERIAL AND METHODS**

### **Sample collection.**

Adult worker bees samples were collected from 13 Collapsing Colonies originating from 7 widely separated apiaries from the Ajloun province. The samples were collected randomly from inside the colonies, transported on ice and transferred to -25 C until use.

### **RNA extraction.**

Individual bees were homogenized in by using pestle and mortar with 1 ml Trizol reagent (Biobasic) and transferred to 1.5 ml microfuge tubes. The protein fraction was separated out in the organic (lower) phase using chloroform and the RNA was precipitated from the aqueous (upper) phase with 1 ml of iso-propanol and centrifugation (14000 rpm). The resultant nucleic acid pellets were washed two times with 70% ethanol and resuspended in nuclease-free water.

### **RT-PCR amplification.**

Each RNA sample was analysed for the presence of bee virus sequences using the Access RT-PCR system (Promega USA) according to the company protocol. Amplification occurred in a 25- $\mu$ l reaction mixture containing 1x reaction buffer, 0.2mM dNTP, 1  $\mu$ M each of the two primers for each virus, 2 mM MgSO<sub>4</sub>, 0.1 U of avian myeloblastosis virus reverse transcriptase, 0.1 U of Tfl DNA polymerase, and 250 to 500 ng of total RNA. The primer sequences and expected length of the RT-PCR products are shown in Table 1. Reverse transcription at 48°C for 48 min was followed by 40 cycles of 95°C for 30 s, 55°C for 1 min, and 68°C for 2 min and a final extension at 68°C for 7 min. Negative controls were included in each RT-PCR experiment. Ten microliters of amplified product was electrophoresed in a 2% agarose gel and stained with ethidium bromide. Bands were photographed under UV light. Fragment sizes were determined with reference to 100-bp ladder (Promega).

**TABLE 1. The sequences of the six primers pairs**


| <b>No.</b> | <b>The Virus</b> | <b>5'-3' Primer*</b>  | <b>Weight (pb)</b> |
|------------|------------------|---|--------------------|
| <b>1</b>   | BQCV             | F: tggtcagctcccactaccttaaac<br>R: gcaacaagaagaaacgtaaaccac    | 700                |
| <b>2</b>   | DWV              | F: cttactctgccgtcgccca<br>R: ccgtaggaactcattatcgcg            | 194                |
| <b>3</b>   | KBV              | F: gatgaacgtcgacctattga<br>R: tgtgggttgctatgagtca             | 415                |
| <b>4</b>   | SBV              | F: gctgaggtaggatctttgcgt<br>R: tcatcatcttcaccatccga           | 824                |
| <b>5</b>   | ABPV             | F: ttatgtgtccagagactgtat<br>R: gctcctattgctcggttttc           | 900                |
| <b>6</b>   | CBPV             | F: agttgtcatggttaacaggatacgag<br>R: tctaactcttagcacgaaagccgag | 455                |

## RESULTS AND DISCUSSION

The Ajlon province is the second most important region in Jordan for beekeeping, in terms of the number of beekeepers and managed colonies. Adult worker bees collected from thirteen colonies in seven widely dispersed apiaries from the Ajlon province were assayed for the presence of six honeybee viruses (ABPV, BQCV, CBPV, DWV, KBV and SBV) using RT-PCR. The results are shown in Figure 1. Out of the 13 colonies examined, 92% were infected with DWV, 8% with SBV and 16% with ABPV. None were infected with CBPV, BQCV or KBV. Several colonies were infected with more than one virus. This indicated that DWV is the most common virus in collapsing bee colonies in Jordan, These data are similar to comparable surveys conducted elsewhere (Tentcheva et al., 2004a). Nearly all the samples from colonies that presented bee mortality were infected with at least one virus or co-infected with more than one virus.

This is the first instance of nucleic acid-based bee virus detection in Jordan. Robinson (1981) previously observed sacbrood virus symptoms in Jordan, and several bee viruses have been previously detected serologically in other countries of the Arabian peninsula (Allen and Ball, 1996). DWV has been detected by RT-PCR in the United Arab Emirates (Berenyi et al., 2007) although no other viruses were surveyed in these studies.

These preliminary results show the presence of several bee viruses, in particular DWV, in dying Jordanian bee colonies. This is similar to surveys conducted in other countries (Nordström et al., 1999; Antunez et al., 2006; Tentcheva et al., 2004b; Berenyi et al., 2006; Todd et al., 2007; Cox-Foster et al., 2007). Further research is needed to determine which other factors are also differentially associated with colony mortality, such as infestation with parasitic mites (*Varroa destructor*, *Acarapis woodi*), *Nosema apis* and *N. cerana* (PSU, 2007), bacterial diseases and any possible effects of chemical treatment of colonies or foraging resources. These results are also important to increase the awareness among beekeepers, bee researchers and the bee-dependent agricultural industries in Jordan and the Arab world that the diseases and parasites that threaten the bee industry world-wide are similarly present in Arab countries, and are similarly associated with bee colony mortality.



**Fig 1:** Detection of different viruses in the Jordanian affected hives. RT-PCR was performed with differential primers amplifying virus-specific sequences. Panel A: Deformed Wing Virus (DWV), Panel B: Sac Brood Virus (SBV), Panel C: Acute Bee Paralysis Virus (ABPV). Lanes 1-13: Different bee samples. Lane 14: Negative control, RT-PCR without a template. Lane M: DNA size marker.

The results of this work were published in a paper at the Jordan Journal of Agricultural Sciences in September 2008-07-12. under the title First detection of honey bee viruses in Jordan by RT-PCR. Based on the results of this work, the Bee Research Unit of Jordan been able to get local and international grants to conduct a research entitled: "Prevalence of honeybee viruses in the Middle East and their association with the appearance of Colony Collapse Disorder (CCD)", Almost certainly this is the first project in the Arab World to suggest using molecular techniques for the detection of bee viruses. The project is conducted in cooperation with Professor Sela's laboratory at the (HUJ). Also the laboratory of the Bee Research Unit is the first in the Arab World to use molecular techniques for the detection of the honey bee viruses. Information on the impact of viral diseases in bees in the Middle East will be gathered. Concurrently, a practical means to control viral diseases and the outcome of CCD will also be investigated. Specific objectives are: 1) Identifying virus diseases of bees in the Middle East, 2) Study the integration of viruses into the bee genome and the engendering of virus resistance. 3) Correlation of CCD with bees that have a history of IAPV infection and the association with Varroa infestation. A project vision is to ultimately be able to breed virus resistant bees. The project is granted by the MERC program of the USAID and it started in January 2009.

## Acknowledgements

The research group would like to thank the Vita-Europe and the National Center for Agricultural Research and Extension for financial support, this project was mainly supported by the **Vita-Europe award 2007**. Special acknowledgments to the original paper coauthors, Mohammad Brake from Jerash Private University- Jordan., Hussein Migdadi from National Center for Agricultural Research and Extension – Jordan, and Joachim R. de Miranda from Queen’s University Belfast Northern Ireland,

## REFERENCES

- **Allen, M.F.**, Ball, B.V. 1996. The incidence and world distribution of honey bee viruses. *Bee World* 77, 141–162.
- **Antunez, K.A.**, D’Alessandro, B., Corbella, E., Ramallo, G., Zunino, P. 2006. Honeybee viruses in Uruguay. *Journal of Invertebrate Pathology* 93: 67–70.
- **Bailey, L.**, 1964. The ‘Isle of Wight disease’: the origins and significance of the myth. *Bee World* 45, 32-37.
- **Bailey, L.**, Ball, B.V. 1991. *Honey Bee Pathology*, second ed. Academic Press, London
- **Bailey, L.**, Ball, B.V., Perry, J.N. 1981. The prevalence of viruses of honey bees in Britain. *Ann. Appl. Biol.* 97, 109-118.
- **Bailey, L.**, Ball, B.V., Perry, J.N. 1983. Association of viruses with two protozoal pathogens of the honey bee. *Ann. Appl. Biol.* 103, 13-20.
- **Bailey, L.**, Perry, J.N. 2001. The natural control of the tracheal mite of honey bees. *Exp. Appl. Acarol.* 25, 745-749.
- **Ball, B.V.**, Bailey, L. 1997. Viruses. In: Morse, R.A., Flottum, K. (Eds.), *Honey Bee Pests Predators and Diseases*. Al Root Co., Medina, pp. 11–31.
- **Berenyi, O.**, Bakonyi, T., Derakhshifar, I., Köglberger, H., Nowotny, N. 2006. Occurrence of six honeybee viruses in diseased Austrian apiaries. *Appl. Environ. Microbiol.* 72, 2414-2420.
- **Berenyi, O.**, Bakonyi, T., Derakhshifar, I., Köglberger, H., Topolska, G., Ritter, W., Pechhacker, H., Nowotny, N. 2007. Phylogenetic analysis of deformed wing virus genotypes from diverse geographic origins indicates recent global distribution of the virus. *Appl. Environ. Microbiol.* 73, 3605-3611.

- **Beuhne, R.**, 1910. Bee mortality. J. Dept. Agric. Victoria 8, 149-151.
- **Chantawannakul, P.**, Ward, L., Boonham, N., Brown, M. 2006. A scientific note on the detection of honeybee viruses using real-time PCR (TaqMan) in *Varroa* mites collected from a Thai honeybee (*Apis mellifera*) apiary. J. Invertebr. Pathol. 91, 69–73.
- **Chen, Y.P.**, Higgins, J.A., Feldlaufer, M.F. 2005. Quantitative analysis of deformed wing virus infection in the honey bee, *Apis mellifera* L. by real-time RT-PCR. Appl. Environ. Microbiol. 71, 436–441.
- **Cox-Foster, D.L.**, Conlan, S., Holmes, E.C., Palacios, G., Evans, J.D., Moran, N.A., Quan, P.-L., Briese, T., Hornig, M., Geiser, D.M., Martinson, V., van Engelsdorp, D., Kalkstein, A.L., Drysdale, A., Hui, J., Zhai, J., Cui, L., Hutchinson, S.K., Simons, J.F., Egholm, M., Pettis, J.S., Lipkin, W.I. 2007. A metagenomic survey of microbes in honey bee colony collapse disorder. Science 318, 283-287.
- **Foote, H.L.**, 1966. The mystery of the disappearing bees. Am. Bee J. 106, 126-127.
- Goodacre, W.A. 1943. Dwindling troubles may cause heavy mortality. Australasian Beekeeper 45, 57-59.
- **Grabensteiner, E.**, Bakonyi, E., Ritter, W., Pechhacker, H., Nowotny, N. 2007. Development of a multiplex RT-PCR for the simultaneous detection of three viruses of the honeybee (*Apis mellifera* L.): Acute bee paralysis virus, Black queen cell virus and Sacbrood virus. Journal of Invertebrate Pathology. 94: 222–225.
- **Haddad, N, Shammout, A, Al-Nsour, A.**, 2007. The economic value of honeybees for crop pollination in Jordan. Documents of the 40th Apimondia International Apicultural Congress, Melbourne, Australia. p115.
- **Hornitzky, M.A.Z.**, 1987. Prevalence of virus infections of honey bees in Eastern Australia. J. Apic. Res. 26, 181-187.
- **Nordstrom, S.**, I. Fries, A. Aarhus, H. Hansen, and S. Korpela. 1999. Virus infection in Nordic honey bee colonies with no, low or severe *Varroa jacobsoni* infestations. Apidologie 30:475–484.
- **Oertel, E.**, 1965. Many bee colonies dead of an unknown cause. Am. Bee J. 105, 48-49.
- PSU. 2007. Bee Researchers Close in on Colony Collapse Disorder. Am. Bee J. 147, 930-931.
- **Rennie, J.**, White, P.B., Harvey, E.J. 1921. Isle of Wight disease in hive bees. Trans. Roy. Soc. Edinb. 52, 737-779.
- **Robinson, W.S.**, 1981. Beekeeping in Jordan. Bee World 62, 91-97.
- **Stokstad, E.**, 2007a. The case of empty hives. Science 316, 970-972.
- **Stokstad, E.**, 2007b. Puzzling decline of U.S. bees linked to virus from Australia. Science 317, 1304-1305.
- **Suchail, S.**, Debrauwer, L., Belzunces, L.P. 2004. Metabolism of imidacloprid in *Apis mellifera*. Pest. Manag. Sci. 60, 291–296.
- **Tentcheva, D.**, Gauthier, L., Jouve, S., Canabady-Rochelle, L., Dainat, B., Cousserans, F., Colin, M.E., Ball, B.V., Bergoin, M. 2004a. Polymerase chain reaction detection of deformed wing virus (DWV) in *Apis mellifera* L., and *Varroa destructor*. Apidologie 35, 431–439.
- **Tentcheva, D.**, Gauthier, L., Zappulla, N., Dainat, B., Cousserans, F., Colin, M. E., Bergoin, M. 2004b. Prevalence and seasonal variations of six bee virus in *Apis mellifera* L. and *Varroa destructor* mite populations in France. Appl. Environ. Microbiol. 70, 7185-7191.
- **Todd, J.H.**, de Miranda, J.R., Ball, B.V. 2007. Incidence and molecular characterization of viruses found in dying New Zealand honey bee (*Apis mellifera*) colonies infested with *Varroa destructor*. Apidologie 38, 354-367.
- **Topley, E.**, Davison, S., Leat, N., Benjeddou, M. 2005. Detection of three honeybee viruses simultaneously by a single multiplex reverse transcriptase PCR. Afric. J. Biotech. 4, 763-767.
- **Wilson, W.T.**, Menapace, D.M. 1979. Disappearing disease of honey bees: A survey of the United States. American Bee Journal 118-119; 184-186; 217.